

## Relation between dynamics of network systems and those of theirs sub-networks <sup>(1)</sup>

Yunjiao Wang<sup>†</sup>, Kiran Chilakamarri<sup>†</sup>, Demetrios Kazakos <sup>†</sup> and Maria C.A. Leite <sup>‡</sup>

 †Department of Mathematics, Texas Southern University Houston, Texas, USA
‡Mathematics & Statistics, University of South Florida at St. Petersburg St. Petersburg, Florida, USA
Email: wangyx@tsu.edu, kazakosd@tsu.edu, mcleite@mail.usf.edu

**Summary.** Biological networks such as gene regulatory networks, neural networks, and metabolic networks are generally complex even from the network topology point of view [17, 18]. However, the understanding of the dynamics of such network systems is crucial to identify mechanisms behind many kinds of biological processes and diseases, and to decode the mysteries of life. Statistical studies on the topology of real world networks revealed some very intriguing features [17] including power-law degree distributions [2, 25, 36], local community structures [4, 11, 13] and network motifs [14, 6]. There is a large body of work devoted to identifying communities or motifs in biological networks [17, 35, 22, 18, 23, 6, 14]. Interestingly, only very few works focused on using modular idea to study dynamics of network systems: the dynamics of a complex network can be understood by studying its subnetwork systems. In order for this idea to work, the dynamics of the subnetworks need to be preserved or partially preserved in the original network. A simple example where this is true is when a subnetwork does not receive input from the rest of the network. However, the situation becomes quite subtle when the subnetwork and its complementary subnetwork have mutual interactions.

In this work, we address the relations between dynamics of the subnetworks and that of the whole system in a Boolean network mathematical modeling framework.

Mathematical models have proven to be indispensable tools for network systems. Among various mathematical modeling frameworks, coupled differential equations and Boolean networks are popular for modeling regulatory networks[7, 26, 16, 20, 21, 3, 10, 33, 1, 12, 29, 30, 31, 15, 5, 28]. Network systems are often represented by directed graphs, wherein components are represented by nodes and interactions by arrows. An n-node Boolean network system is a discrete dynamical system with the form X(t+1) =F(X(t)), where  $X = (x_1, \dots, x_n)$ ,  $x_i$  represents the state variable of the  $i^{th}$  node,  $F = (f_1, \dots, f_n)$  and  $f_i$  is the governing function of the  $i^{th}$  node with its value being either 0 or 1. Boolean networks have been widely used to model biological regulatory networks [7, 26, 16, 20, 21, 3, 10, 33, 1, 12]. They can be set up in situations where information on the detailed kinetic interactions is not available and can provide valuable insights [19, 12, 24, 27, 8, 9, 32].

In this work, we particularly consider networks formed by two subnetworks connected at a cutting node, which we will define next. A node is called a *cutting node* of a connected network if the removal of the node leads to two or more disjoint subnetworks. Furthermore, we introduce the notion of a network being agreeable. Let G be the network of the whole system formed by  $G_1$  and  $G_2$  connected at a cutting node c. Let  $x_c(t, *)$  be the value of the cutting node in the system \* (here \* can be  $G_1$ , G, or  $G_2$ ) at time t. We say that G is agreeable if  $x_c(t,G) = z_0$  whenever  $x_c(t,G_1) = x_c(t,G_2) = z_0$ . Additionally, we give an example of an updating scheme for the cutting node that guarantees that a network system is agreeable. We show that if a network is agreeable and its subnetworks have only cycles, then the whole system has only cycles. We then prove that if  $X_0$  is a fixed point of G, then  $X_0$  restricted to the phasespace of one of the subnetwork systems must be a fixed point of that system.

We also discuss the relations between the product of the transition diagrams (a representation of trajectories) of the subnetwork systems and that of the whole system. In a Boolean network system, the transition diagram of the system represents the dynamics of the system. When the dynamics of the subnetworks are all independent, then the dynamics of the whole network is just the product of the subnetworks. However, when they are not independent, the relation is not all that transparent. We discuss the construction of the transition diagram of a network G from the transition diagram of the product system of its subnetworks and present an algorithm to construct the transition diagram of G. This algorithm can be very useful when the subnetworks are large and their transition diagrams are ready to use. On the other hand, the algorithm provide a rather clear view on the relations between the dynamics of the whole network and that of its subnetworks.

Acknowledgments. A few days before KC passed away, he discussed with YW about finishing this work in the summer of 2015. YW has been grateful for KCs friendship and mentoring. YW was supported by DHS-14ST-062-001.

<sup>(1)</sup> This work has been published in AIMS Mathematics [34]

## References

- W. ABUO-JAOUDE, D. OUATTARA, AND M. KAUFMAN, From structure to dynamics: frequency tuning in the p53-mdm2 network: I. logical approach, Journal of Theoretical Biology, 258 (2009), pp. 561–577.
- [2] R. Albert, Scale-free networks in cell biology, Journal of Cell Science, 118 (2005), pp. 4947–4957.
- [3] R. ALBERT AND H. OTHMER, The topology of the regulatory interactions perdicts the expression pattern of the segment polarity genes in drosophila melanogaster, Journal of Theoretical Biology, 233 (2003), pp. 1–18.
- [4] S. A. ALCALÁ-CORONA, T. E. VELÁZQUEZ-CALDELAS, J. ESPINAL-ENRÍQUEZ, AND E. HERNÁNDEZ-LEMUS, Community structure reveals biologically functional modules in mef2c transcriptional regulatory network, Front Physiol., 7 (2016).
- [5] B. B. ALDRIGE, J. M. BURKE, D. A. LAUFFENBURGE, AND P. SORGER, *Physicochemical modeling of cell signaling pathways*, Nature Cell Biol, 8 (2006).
- [6] U. ALON, Network motifs: Theory and experimental approaches, Nature Reviews Genetics, 8 (2007), p. 450.
- [7] C. CAMPBELL, J. THAKAR, AND R. ALBERT, Network analysis reveals cross-links of the immune pathways activated by bacteria and allergen, Physical Reveiw. E, Statistical physics, plasmas, fluids and related interdisciplinary topics, 84 (2011), p. 031929.
- [8] R. EDWARDS AND L. GLASS, Combinatorial explosion in model gene networks, Chaos: An inerdisciplinary Journal of Nonlinear Science, 11 (2000), pp. 691– 704.
- [9] R. EDWARDS, H. SIEGELMANN, K. AZIZA, AND L. GLASS, Symbolic dynamics and computation in model gene models, Chaos: An inerdisciplinary Journal of Nonlinear Science, 11 (2001), pp. 160–169.
- [10] C. ESPIONZA-SOTO, P. PADILLA-LONGORIA, AND E. ALVAREZ-BUYLLA, A gene regulatory network model for cell-fate determination during arabidopsis thaliana flower development that is robus and recovers experiental gene expression profiles, Plant Cell, 16 (2004), pp. 2923–2939.
- [11] N. M. E. GIRVAN M., Community structure in social and biological networks, Proc. Natl. Acad. Sci. U.S.A., 99 (2002), pp. 7821–7826.
- [12] L. GLASS AND S. A. KAUFFMAN, The logical analysis of continuous, nonlinear biochemical control network, J. Theor. Biol., 39 (1973), pp. 103–139.

- [13] C. L., M. E., F. S., AND C. M, Detection of gene communities in multi-networks reveals cancer drivers., Sci. Rep., 5 (2015).
- [14] R. MILO, S. SHEN-ORR, S. ITZKOVITZ, N. KASHTAN, D. CHKLOVSKII, AND U. ALON, *Network motifs: Simple building blocks of complex networks*, Science, 298 (2002), pp. 824–827.
- [15] A. MOGILNER, R. WOLLMAN, AND W. F. MARSHALL, *Quantitative modeling in cell biology*, Developmental Cell, 11 (2006), pp. 279–287.
- [16] S. L. ND S. M. ASSMANN AND R. ALBERT, Predicting essential components of signal transduction networks: A dynamic model of guard cell abscisic acid signaling, PLoS Biol., 4 (2006), p. e312.
- [17] M. E. J. NEWMAN, Modularity and community structure in networks, Proceedings of the National Academy of Sciences, 103 (2006), pp. 8577–8582.
- [18] M. E. J. NEWMAN, Communities, modules and large scale structure, Nature Physics, 8 (2012), pp. 25–31.
- [19] A. SAADATPOUR, R. ALBERT, AND T. C. RELUGA, A reduction method for boolean network models proven to conserve attractors, SIAM J. Appl. Dyn. Sys., 12 (2013), pp. 1997–2011.
- [20] J. SAEZ-RODRIGUEZ, L. SIMEONI, J. A. LINDQUIST, R. HEMENWAY, U. BOMMHARDT, U. U. H. B. ARNDT, R. WEISMANTEL, E. D. GILLES, S. KLAMT, AND B. SCHRAVEN, A logical model provides insights into t cell receptor signaling, PLoS Computational Biology, 3 (2007), p. e163.
- [21] L. SANCHEZ AND D. THIEFFRY, A logical analysis of the drosophila gap-gene system, J. Theor. Biol., 211 (2001), pp. 115–141.
- [22] A. J. SCHWARZ, A. GOZZI, AND A. BIFONE, *Community* structure and modularity in networks of correlated brain activity, Resonance Imaging.
- [23] S. S. SHEN-ORR, R. MILO, S. MANGAN, AND U. ALON, Network motifs in the transcriptional regulation network of escherichia coli, Nature Genet, 31 (2002), pp. 64–68.
- [24] E. SNOUSSI, Qualitative dynamics of piecewise differential equations: a discrete mapping approach., Dynamical and Stability of Systems, 4 (1989), pp. 189– 207.
- [25] R. TANAKA, *Scale-rich metabolic networks.*, Phys. Rev. Lett., 94 (2005), p. 168101.
- [26] J. THAKAR, A. K. PATHAK, L. MURPHY, R. ALBERT, I. CATTADORI, AND R. J. D. BOER, Network model of immune responses reveals key effectors to single and

*co-infection dynamics by a respiratory bacterium and a gatrointestinal helminth*, PLoS Computational Biology, 8 (2012).

- [27] R. THOMAS, Biological Feedback, CRC, 1990.
- [28] D. TURNER, P. PASZEK, D. WOODCOCK, D. E. NEL-SON, C.A.HORTON, Y. WANG, D. SPILLER, D. A. RAND, M. R. H. WHITE, AND C. V. HARPER, *Physiological levels of tnfalpha stimulation induce stochastic dynamics of nf-kappab responses in single living cells*, Journal of Cell Biology, 123 (2010), pp. 2834–2843.
- [29] J. TYSON AND B. NOVAK, Functional motifs in biochemical reaction networks, Annu. Rev. Phys. Chem., 61 (2010).
- [30] J. J. TYSON, K. C. CHEN, AND B. NOVAK, Network dynamics and cell physiology, Nature Rev. Mol. Cell Biol, 2 (2001), pp. 908–916.
- [31] ——, Sniffers, buzzers, toggles and blinkers: dynamics of regulatory and signaling pathways in the cell, Curr. Op. Cell Biol, 15 (2003).
- [32] A. VELIZ-CUBA, A. KUMAR, AND K. JOSIC, Piecewise linear and boolean models of chemical reaction networks, accepted to J. Math. Bio., (2014).
- [33] R. S. WANG AND R. ALBERT, Discrete dynamical modeling of cellular signaling networks, Methods in Enzymology, 467 (2009), pp. 281–306.
- [34] Y. WANG, K. CHILAKAMARRI, D. KAZAKOS, AND M. LEITE, *Relations between the dynamics of network* systems and their dynamics, AIMS Mathematics, 2(3) (2017), pp. 437–450.
- [35] S. WERNICKE, A faster algorithm for detecting network *motifs*, Proc. 5th WABI-05, 3692 (2005).
- [36] S. H. YOOK, Z. N. OLTVAI, AND A. L. BARABÁSI, Functional and topological characterization of protein interaction networks, Proteomics, 4 (2004), pp. 928– 942.