

Intrinsic Localized Modes in a Magnetically Coupled Two-degree-of-freedom Resonator Array

Masayuki Kimura[†]

 †Faculty of Science and Engineering, Setsunan University
17–8 Ikeda-Nakamachi, Neyagawa, Osaka 572-8508, Japan Email: masayuki.kimura@setsunan.ac.jp

Summary

Nonlinearly coupled resonator array is known to have spatially localized and temporally periodic solutions which are called intrinsic localized modes (ILMs) or discrete breathers (DBs)[1]. It is well known that the ILM/DB exist not only in one-dimensional array but also in two- or three-dimensional lattice[2]. Many studies of ILM/DB for one-dimensional arrays assume the array is rigid *i.e.* each oscillator moves only along the axis of the array.

A flexible nonlinear chain in which each oscillator can move two-dimensionally is known to have two types of intrinsic localized mode(ILM). One is the same as the traditional ILM that oscillates along the chain axis. We call it longitudinal ILM (L-ILM). The other mainly oscillates perpendicularly, which is called transverse ILM (T-ILM). In a flexible Fermi–Pasta–Ulam chain which has cubic nonlinearity in the coupling force, L-ILM loses its stability because of the buckling effect of the chain[3]. For T-ILM, almost all the solutions are unstable when the chain is not stretched[4]. If the chain has nonlinearity in the on-site restoring force, stable L- and T-ILM can be found for certain regions in the ω - β plane where ω is the angular frequency of ILM and β is the ratio of on-site and inter-site nonlinearity[5].

A magnetically coupled two-degree-of-freedom resonator array is designed for experimental studies for L- and T-ILM in a flexible chain. An elastic rod with a cylindrical permanent magnet at the tip is a two-degree-of-freedom oscillator. The permanent magnet induces nonlinearity in the coupling force of adjacent oscillators. In this study, the magnets are attached to generate repulsive force and it becomes stronger if the distance between adjacent oscillators becomes closer. Then the nonlinearity of the coupling force is classified as hard spring nonlinearity. ILM will appear above the upper bound of the phonon band. If the edge of the array is sinusoidally excited by an external exciter at a frequency higher than the upper bound, several moving L-ILMs appear and travel along the chain[6, 7]. The moving L-ILMs can keep their shape of localized energy distribution unless the excitation amplitude is rather small. However, if the amplitude becomes large, transverse oscillations are induced and the moving L-ILMs become



Figure 1: Magnetically coupled two-degree-of-freedom resonator array

unstable. The instability will come from the flexibility of the chain. There is no force to prevent the deformation of the flexible chain except the restoring force of the rod. Additional permanent magnet arrays are placed parallel to the flexible chain. Nonlinearity is added to the restoring force against the transverse displacement. In this report, the stability of static L- and T-ILMs is investigated for the flexible chains. We will discuss the effect of the surrounding resonator arrays on the stability of ILMs.

Acknowledgments

We are grateful to Prof. Shnji Doi and Mr. Lee Jung-Jin for helpful discussions on the model of the magnetically coupled 2D chain. We also would like to thank Prof. Alain Bertrand Togueu Motcheyo for fruitful discussion on the supratransmission phenomenon which is related to this project. This work is partially supported by JSPS Kakenhi No. 18K04020 and No. 21K03935.

References

- [1] A.J. Sievers and S. Takeno, Phys. Rev. Lett., vol.61, pp.970, 1988.
- [2] S. Flach and A. V. Gorbach, Phys. Rep., vol.467, pp.1–116, 2008.
- [3] M. Kimura, A. Mitani, and S. Doi, *Lett. on Mater.*, vol.6(1), pp.22–26, 2016.
- [4] M. Kimura, A. Mitani, and S. Doi, *Nonlinear Theory and Its Applications, IEICE*, vol.8(2), pp.153–161, 2017.
- [5] I. Yamaguchi, M. Kimura, and S. Doi. "Finding moving intrinsic localized modes based on stability of static intrinsic localized modes in flexible FPU-KG mixed lattices", In *NOLTA2020*, Online, Japan, November 16-19, 2020.
- [6] M. Kimura, J.-J. Lee, and S. Doi, *Tech. Rep. of IEICE*, vol.121(61), pp.5–9, 2021.
- [7] M. Kimura, J.-J. Lee, A.B. Togueu Motcheyo, and S. Doi. "Nonlinear supratransmission in a magnetically coupled elastic rods arranged in three lines", In *Book of Abstracts: LONE 2022 in CMD29*, Manchester, United Kingdom / Hybrid, August 21-26, 2022.

ORCID iDs Masayuki Kimura: 10 0000-0002-1445-6266



This work is licensed under a Creative Commons Attribution NonCommercial, No Derivatives 4.0 License.